



Principles of Home Canning



**Featuring new, research-based
recommendations for safer and
better quality food at home**

Know Your Altitude

It is important to know your approximate elevation or altitude above sea level in order to determine a safe processing time for canned foods. Since the boiling temperature of liquid is lower at higher elevations, it is critical that additional time be given for the safe processing of foods at altitudes above sea level.

All towns and communities in Kentucky are below 2,000 feet. The processing times given in this canning guide are for altitudes up to 3,000 feet and are safe for all parts of our state.

DANGER!

Guard Against Food Poisoning

Pressure canning is the only recommended method for canning meat, poultry, seafood, and most vegetables. The bacterium *Clostridium botulinum* in low-acid foods is destroyed when they are processed at the correct time and pressure in pressure canners. Using boiling-water canners for these foods poses a real risk of botulism poisoning.

If these bacteria survive and grow inside a sealed jar of food, they can produce a poisonous toxin. Even a taste of food containing this toxin can be fatal. Low-acid foods should be boiled after their jars are opened, even if you detect no signs of spoilage and are certain the food has been properly processed. In Kentucky, boiling for 13 minutes destroys the toxin that causes poisoning.

This guide, containing up-to-date instructions for preparing safe home-canned foods, is based on research conducted at the Pennsylvania State University. It is adapted from the USDA Complete Guide to Home Canning.

Other publications in the home-canning series include:

- FCS3-121, *Canning Tomatoes and Tomato Juice*
- FCS3-326, *Selecting, Preparing, and Canning Fruit and Fruit Products*
- FCS3-327, *Selecting, Preparing, and Canning Tomatoes and Tomato Products*
- FCS3-328, *Selecting, Preparing, and Canning Vegetables and Vegetable Products*
- FCS3-329, *Preparing and Canning Poultry, Red Meat, and Fish*
- FCS3-330, *Preparing and Canning Fermented Foods and Pickled Vegetables*
- FCS3-331, *Preparing and Canning Jams and Jellies*

These publications are also available on our web site at:

<http://www.ca.uky.edu/agc/pubs/pubs.htm>

No product endorsement is implied, nor is discrimination against similar products intended, by the mention of brand names in this publication.

Revised and adapted for use in Kentucky by Sue Burrier, former Extension Food and Nutrition Specialist

*Contact: Sandra Bastin, Ph.D., R.D.,
Extension Specialist in Food and Nutrition*

Contents

Why Can Foods?	4
How Canning Preserves Foods	4
Ensuring Safe Canned Foods.....	4
Food acidity & processing methods ...	5
Equipment & methods NOT recommended	5
Ensuring High-Quality Canned Foods	6
Maintaining color & flavor in canned food	6
Advantages of hot packing	7
Controlling headspace	7
Jars and Lids	7
Jar cleaning	8
Sterilization of empty jars	8
Lid selection, preparation & use	8
Selecting the Correct Processing Time ..	9
Recommended Canners	9
Boiling-water canners	9
Using boiling-water canners	10
Pressure canners	10
Using pressure canners	11
Cooling Jars	11
Testing Jar Seals	12
Reprocessing Unsealed Jars	12
Storing Canned Foods	12
Identifying and Handling Spoiled	
Canned Food	12
Detoxification process	13
Canned Foods for Special Diets	13
Canning without sugar	13
Canning without salt <i>(reduced sodium)</i>	13
Glossary	14
Do Your Canned Foods Pass	
This Quality Test?	16

Introduction to the Complete Guide to Home Canning

Home canning has changed greatly in the 170 years since it was introduced as a way to preserve food. Scientists have found ways to produce safer, higher quality products. The first part of this series explains the scientific principles on which canning techniques are based, discusses canning equipment, and describes the proper use of jars and lids. It describes basic canning ingredients and procedures and how to use them to achieve safe, high-quality canned products.

The remaining publications in this series consist of canning guides for specific foods. These guides offer detailed directions for making sugar syrups and for canning fruits and fruit products, tomatoes and tomato products, vegetables, red meats, poultry, seafoods, pickles, and relishes. Handy guidelines for choosing the correct quantity and quality of raw foods accompany each set of directions for fruits, tomatoes, and vegetables. Most recipes are designed to yield a full canner load of pints or quarts.

This publication contains many new research-based recommendations for canning safer and better quality food at home. It is an invaluable resource for persons who are canning for the first time. Experienced canners will find updated information to help them improve their canning practices.

Why Can Foods?

Canning can be a safe and economical way to preserve quality food at home. Disregarding the value of your labor, canning homegrown food may save you half the cost of buying commercially canned food. Canning favorite and special products to be enjoyed by family and friends is a fulfilling experience and a source of pride for many people.

Many vegetables begin losing some of their vitamins when harvested. Nearly half the vitamins may be lost within a few days unless the fresh produce is cooled or preserved. Within one to two weeks, even refrigerated produce loses half or more of some of its vitamins. The heating process during canning destroys from one-third to one-half of vitamins A and C, thiamin, and riboflavin. Once canned, additional losses of these sensitive vitamins are from 5 percent to 20 percent each year. The amounts of other vitamins, however, are only slightly lower in canned compared with fresh food. If vegetables are handled properly and canned promptly after harvest, they can be more nutritious than fresh produce sold in local stores.

The advantages of home canning are lost when you start with poor quality fresh foods, when jars fail to seal properly, when food spoils, and when flavors, texture, color, and nutrients deteriorate during prolonged storage.

The information and guides that follow explain many of these problems and recommend ways to minimize them.

How Canning Preserves Foods

The high percentage of water in most fresh foods makes them very perishable. They spoil or lose their quality for several reasons:

- growth of undesirable microorganisms—bacteria, molds, and yeasts,

- activity of food enzymes,
- reactions with oxygen, and
- moisture loss.

Microorganisms live and multiply quickly on the surfaces of fresh food and on the inside of bruised, insect-damaged, and diseased food. Oxygen and enzymes are present throughout fresh food tissues.

Proper canning practices include:

- carefully selecting and washing fresh food,
- peeling some fresh foods,
- hot packing many foods,
- adding acids (lemon juice, citric acid, or vinegar) to some foods,
- using acceptable jars and self-sealing lids, and
- processing jars in a boiling-water or pressure canner for the correct time.

Collectively, these practices remove oxygen; destroy enzymes; prevent the growth of undesirable bacteria, yeasts, and molds; and help form a high vacuum in jars. High vacuums form tight seals which keep liquid in and air and microorganisms out.

Ensuring Safe Canned Foods

Growth of the bacterium *Clostridium botulinum* in canned food may cause botulism—a deadly form of food poisoning. These bacteria exist either as spores or as vegetative cells. The spores, which are comparable to plant seeds, can survive harmlessly in soil and water for many years. When ideal conditions exist for growth, the spores produce vegetative cells which multiply rapidly and may produce a deadly toxin within three to four days of growth in an environment consisting of:

- a moist, low-acid food,
- a temperature between 40°F and 120°F, and
- less than 2 percent oxygen.

Botulinum spores are on most fresh food surfaces. Because they grow only in the absence of air, they are harmless on fresh foods.

Most bacteria, yeasts, and molds are difficult to remove from food surfaces. Washing fresh food reduces their numbers only slightly. Peeling root crops, underground stem crops, and tomatoes reduces their numbers greatly. Blanching also helps, but the vital controls are the method of canning and use of the recommended research-based processing times found in the publications of this home-canning series. These processing times ensure destruction of the largest expected number of heat-resistant microorganisms in home-canned foods.

Properly sterilized canned food will be free of spoilage if lids seal and jars are stored below 95°F. Storing jars at 50°F to 70°F enhances retention of quality.

Food acidity & processing methods

Whether food should be processed in a pressure canner or boiling-water canner to control botulism bacteria depends on the acidity in the food. Acidity may be natural, as in most fruits, or added, as in pickled food. Low-acid canned foods contain too little acidity to prevent the growth of these bacteria. Acid foods contain enough acidity to block their growth or to destroy them rapidly when heated. The term “pH” is a measure of acidity; the lower its value, the more acidic the food. The acidity level in foods can be increased by adding lemon juice, citric acid, or vinegar.

Low-acid foods have pH values higher than 4.6. They include red meats, seafood, poultry, milk, and all fresh vegetables except for most tomatoes. Most products that are mixtures of low-acid and acid foods also have pH values above 4.6 unless their ingredients include enough lemon juice, citric acid, or vinegar to make them acid foods. Acid foods have a pH of 4.6 or lower. They include fruits, pickles, sauerkraut, jams, jellies, marmalade, and fruit butters.

Although tomatoes usually are considered an acid food, some are now known to have pH values slightly above 4.6. Figs also have pH values slightly above 4.6. Therefore, if they are to be canned as acid foods, these products must be acidified to a pH of 4.6 or lower with lemon juice or citric acid. Properly acidified tomatoes and figs are acid foods and can be safely processed in a boiling-water canner.

Botulinum spores are very hard to destroy at boiling-water temperatures; the higher the canner temperature, the more easily they are destroyed. Therefore, all low-acid foods should be sterilized at temperatures of 240°F to 250°F, attainable with pressure canners operated at 10 to 15 PSIG. (PSIG means pounds per square inch of pressure as measured by a gauge.) At these temperatures, the time needed to destroy bacteria in low-acid canned foods ranges from 20 to 100 minutes. The exact time depends on the kind of food being canned, the way it is packed into jars, and the size of jars. The time needed to safely process low-acid foods in boiling water ranges from seven to 11 hours; the time needed to process acid foods in boiling water varies from five to 85 minutes.

Equipment & methods NOT recommended

Open-kettle canning and the processing of freshly filled jars in conventional ovens, microwave ovens, and dishwashers are not recommended because these practices do not prevent all risks of spoilage. Steam canners are not recommended because processing times for use with current models have not been adequately researched. Because steam canners may not heat foods in the same manner as boiling-water canners, their use with boiling-water processing times may result in spoilage. So-called canning powders are useless as preservatives and do not replace the need for proper heat processing.

It is not recommended that pressures in excess of 15 PSIG be applied when using new pressure-canning equipment.

Jars with wire bails and glass caps make attractive antiques or storage containers for dry food ingredients but are not recommended for canning. One-piece, porcelain-lined, zinc caps are also no longer recommended. Both glass and zinc caps use flat rubber rings for sealing jars but too often fail to seal properly.

Ensuring High-Quality Canned Foods

Begin with good-quality fresh foods suitable for canning. Quality varies among varieties of fruits and vegetables. Many county Extension offices can recommend varieties best suited for canning.

Examine food carefully for freshness and wholesomeness. Discard diseased and moldy food. Trim small diseased lesions or spots from food.

Can fruits and vegetables picked from your garden or purchased from nearby producers when the products are at their peak of quality—within six to 12 hours after harvest for most vegetables. For best quality, apricots, nectarines, peaches, pears, and plums should be ripened one or more days between harvest and canning. If you must delay the canning of other fresh produce, keep it in a shady, cool place.

Fresh home-slaughtered red meats and poultry should be chilled and canned without delay. Do not can meat from sickly or diseased animals. Put fish and seafoods on ice after harvest, eviscerate immediately, and can them within two days.

Maintaining color & flavor in canned food

To maintain good natural color and flavor in stored canned food, you must:

- remove oxygen from food tissues and jars,
- quickly destroy the food enzymes, and
- obtain high jar vacuums and airtight jar seals.

Follow these guidelines to ensure that your canned foods retain optimal colors and flavors during processing and storage:

- Use only high-quality foods that are at the proper maturity and are free of diseases and bruises.
- Use the hot-pack method, especially with acid foods to be processed in boiling water.
- Don't unnecessarily expose prepared foods to air; can them as soon as possible.
- While preparing a canner load of jars, keep peeled, halved, quartered, sliced or diced apples, apricots, nectarines, peaches, and pears in a solution of 3 grams (3000 milligrams) ascorbic acid to 1 gallon of cold water. This procedure is also useful in maintaining the natural color of mushrooms and potatoes and for preventing stem-end discoloration in cherries and grapes. You can get ascorbic acid in several forms:

Pure powdered form—seasonally available among canning supplies in supermarkets. One level teaspoon of pure powder weighs about 3 grams. Use 1 teaspoon per gallon of water as a treatment solution.

Vitamin C tablets—economical and available year-round in many stores. Buy 500-milligram tablets; crush and dissolve six tablets per gallon of water as a treatment solution.

Commercially prepared mixes of ascorbic and citric acid—seasonally available among canning supplies in supermarkets. Sometimes citric acid powder is sold in supermarkets, but it is less effective in controlling discoloration. If you choose to use these products, follow the manufacturer's directions.

- Fill hot foods into jars and adjust headspace as specified in recipes.
- Tighten screw bands securely, but if you are especially strong, not as tightly as possible.
- Process and cool jars.
- Store the jars in a relatively cool, dark place, preferably between 50°F and 70°F.
- Can no more food than you will use within a year.

Advantages of hot packing

Many fresh foods contain from 10 percent to more than 30 percent air. How long canned food retains high quality depends on how much air is removed from food before jars are sealed. The more air that is removed, the higher the quality of the canned product.

Raw-packing is the practice of filling jars tightly with freshly prepared but unheated food. Such foods, especially fruit, will float in the jars. The entrapped air in and around the food may cause discoloration within two to three months of storage. Raw-packing is more suitable for vegetables processed in a pressure canner.

Hot-packing is the practice of heating freshly prepared food to boiling, simmering it three to five minutes, and promptly filling jars loosely with the boiled food.

Hot-packing is the best way to remove air and is the preferred pack style for foods processed in a boiling-water canner. At first, the color of hot-packed foods may appear no better than that of raw-packed foods, but within a short storage period both color and flavor of hot-packed foods will be superior.

Whether food has been hot-packed or raw-packed, the juice, syrup, or water to be added to the foods should be heated to boiling before it is added to the jars. This practice helps to remove air from food tissues, shrinks food, helps keep the food from floating in the jars, increases vacuum in sealed jars, and improves shelf life. Preshrinking food permits adding more food into each jar.

Controlling headspace

The unfilled space above the food in a jar and below its lid is termed headspace. Directions for canning specify leaving ¼-inch for jams and jellies, ½-inch for fruits and tomatoes to be processed in boiling water, and from 1 to 1¼ inches in low-acid foods to be processed in a pressure canner.

This space is needed for expansion of food as jars are processed and for forming vacuums in cooled jars. The extent of expansion is determined by the air content in the food and by the processing temperature. Air expands greatly when heated to high temperatures; the higher the temperature, the greater the expansion. Foods expand less than air when heated.

Jars and Lids

Food may be canned in glass jars or metal containers. Metal containers can be used only once. They require special sealing equipment and are much more costly than jars.

It is recommended that Mason-type jars designed for home canning be used for preserving food by pressure or boiling-water canning. Regular and widemouthed threaded Mason jars with self-sealing lids are the best choices. They are available in half-pint, pint, 1.5-pint, and quart sizes. The standard jar mouth opening is about 2³/₈ inches. Widemouthed jars have openings of about 3 inches, making them more easily filled and emptied. Regular-mouth decorator jelly jars are available in 8- and 12-ounce sizes.

With careful use and handling, Mason jars may be reused many times, requiring only new lids each time. When lids are used properly, jar seals and vacuums are excellent.

Jar cleaning

Before reuse, wash empty jars in hot water with detergent and rinse well by hand, or wash in a dishwasher. Unrinsed detergents may cause unnatural flavors and colors. These washing methods do not sterilize jars. Scale or hard-water films on jars are easily removed by soaking jars several hours in a solution containing 1 cup of vinegar (5 percent) per gallon of water.

Sterilization of empty jars

Use sterile jars for all jams, jellies, and pickled products processed less than 10 minutes. To sterilize empty jars, put them right side up on the rack in a boiling-water canner. Fill the canner and jars with hot (not boiling) water to 1 inch above the tops of the jars. Boil 10 minutes. Remove and drain hot sterilized jars one at a time. Save the hot water for processing filled jars. Fill jars with food, add lids, and tighten screw bands.

Empty jars used for vegetables, meats, and fruits to be processed in a pressure canner need not be sterilized beforehand. It is also unnecessary to sterilize jars for fruits, tomatoes, and pickled or fermented foods that will be processed 10 minutes or longer in a boiling-water canner.

Lid selection, preparation & use

The common self-sealing lid consists of a flat metal lid held in place by a metal screw band during processing. The flat lid is crimped around its bottom edge to form a trough, which is filled with a colored gasket material. When jars are processed, the lid gasket softens and flows slightly to cover the jar-sealing surface, yet allows air to escape from the jar. The gasket then forms an airtight seal as the jar cools. Gaskets in unused lids work well for at least five years from date of manufacture. The gasket material in older unused lids may fail to seal on jars.

It is best to buy only the quantity of lids you will use in a year. To ensure a good seal, carefully follow the manufacturer's directions in preparing lids for use. Examine all metal lids carefully. Do not use old, dented, or deformed lids or lids with gaps or other defects in the sealing gasket.

After filling jars with food, release air bubbles by inserting a flat plastic (not metal) spatula between the food and the jar. Slowly turn the jar and move the spatula up and down to allow air bubbles to escape. Adjust the headspace and then clean the jar rim (sealing surface) with a dampened paper towel. Place the lid, gasket down, onto the cleaned jar-sealing surface. Uncleaned jar-sealing surfaces may cause seal failures.

Then fit the metal screw band over the flat lid. Follow the manufacturer's guidelines enclosed with or on the box for tightening the jar lids properly.

- If screw bands are too tight, air cannot vent during processing, and food will discolor during storage. Overtightening also may cause lids to buckle and jars to break, especially with raw-packed, pressure-processed food.
- If screw bands are too loose, liquid may escape from jars during processing, seals may fail, and the food will need to be reprocessed.

Do not retighten lids after processing jars. As jars cool, the contents in the jar contract, pulling the self-sealing lid firmly against the jar to form a high vacuum.

Screw bands are not needed on stored jars. They can be removed easily after jars are cooled. When removed, washed, dried, and stored in a dry area, screw bands may be used many times. If left on stored jars, they become difficult to remove, often rust, and may not work properly again.

Selecting the Correct Processing Time

When food is canned in boiling water, more processing time is needed for most raw-packed foods and for quart jars than is needed for hot-packed foods and pint jars.

To destroy microorganisms in acid foods processed in a boiling-water canner, you must:

- process jars for the correct number of minutes in boiling water.
- cool the jars at room temperature.

To destroy microorganisms in low-acid foods processed with a pressure canner, you must:

- process the jars for the correct number of minutes at 240°F (10 PSIG) or 250°F (15 PSIG).
- allow canner to cool at room temperature until it is completely depressurized.

The food may spoil if you fail to use the proper processing times, fail to vent steam from canners properly, process at lower pressure than specified, process for fewer minutes than specified, or cool the canner with water.

Processing times for half-pint and pint jars are the same, as are times for 1.5-pint and quart jars. For some products, you have a choice of processing at 5, 10, or 15 PSIG. In these cases, choose the canner pressure (PSIG) you wish to use and match it with your pack style (raw or hot) and jar size to find the correct processing time.

Recommended Canners

There are two main types of canners for heat-processing home-canned food: boiling-water canners and pressure canners. Most are designed to hold seven

quart-jars or eight to nine pint-jars. Small pressure canners hold four quart-jars; some large pressure canners hold 18 pint-jars in two layers but hold only seven quart-jars. Pressure saucepans with smaller volume capacities are not recommended for use in canning. Treat small pressure canners as standard larger canners. They should be vented using the typical venting procedures.

Low-acid foods must be processed in a pressure canner to be free of botulism risks. Although pressure canners also may be used for processing acid foods, boiling-water canners are recommended because they are faster. A pressure canner would require from 55 to 100 minutes to can a load of jars; the total time for canning most acid foods in boiling water varies from 25 to 60 minutes.

A boiling-water canner loaded with filled jars requires about 20 to 30 minutes of heating before its water begins to boil. A loaded pressure canner requires about 12 to 15 minutes of heating before it begins to vent, another 10 minutes to vent the canner, another 5 minutes to pressurize the canner, another 8 to 10 minutes to process the acid food, and, finally, another 20 to 60 minutes to cool the canner before removing jars.

Boiling-water canners

These canners are made of aluminum or porcelain-covered steel. They have removable perforated racks and fitted lids. The canner must be deep enough so that at least 1 inch of briskly boiling water will cover the tops of jars during processing. Some boiling-water canners do not have flat bottoms. A flat bottom must be used on an electric range. Either a flat or ridged bottom can be used on a gas burner. To ensure uniform processing of all jars with an electric range, the canner should be no more than 4 inches wider in diameter than the element on which it is heated.

Using boiling-water canners

Follow these steps for successful boiling-water canning:

1. Fill the canner halfway with water.
2. Preheat water to 140°F for raw-packed foods and to 180°F for hot-packed foods.
3. Load filled jars, fitted with lids, into the canner rack and use the handles to lower the rack into the water; or fill the canner, one jar at a time, with a jar lifter.
4. Add more boiling water, if needed, so the water level is at least 1 inch above jar tops.
5. Turn heat to its highest position until water boils vigorously.
6. Set a timer for the minutes required for processing the food.
7. Cover with the canner lid and lower the heat setting to maintain a gentle boil throughout the processing time.
8. Add more boiling water, if needed, to keep the water level above the jars.
9. When jars have been boiled for the recommended time, turn off the heat and remove the canner lid.
10. Using a jar lifter, remove the jars and place them on a towel, leaving at least 1 inch of space between the jars during cooling.

Pressure canners

Pressure canners for use in the home have been extensively redesigned in recent years. Models made before the 1970s were heavy-walled kettles with clamp-on or turn-on lids. They were fitted with a dial gauge, a vent port in the form of a petcock or counterweight, and a safety fuse. Modern pressure canners are lightweight, thin-walled kettles; most have turn-on lids. They have a jar rack, gasket, dial or weighted gauge, an automatic vent or cover lock, a vent port (steam vent) that is closed with a counterweight or weighted gauge, and a safety fuse.

Pressure does not destroy microorganisms, but high temperatures applied for a

certain period of time do. The success of destroying all microorganisms capable of growing in canned food is based on the temperature obtained in pure steam, free of air, at sea level. At sea level, a canner operated at a gauge pressure of 10 pounds provides an internal temperature of 240°F.

Air trapped in a canner lowers the inside temperature and results in underprocessing. The highest volume of air trapped in a canner occurs in processing raw-packed foods in dial-gauge canners. These canners do not vent air during processing. To be safe, all types of pressure canners must be vented 10 minutes before they are pressurized.

To vent a canner, leave the vent port uncovered on newer models or manually open petcocks on some older models. Heating the filled canner with its lid locked into place boils water and generates steam that escapes through the petcock or vent port. When steam first escapes, set a timer for 10 minutes. After venting 10 minutes, close the petcock or place the counterweight or weighted gauge over the vent port to pressurize the canner.

Weighted-gauge models exhaust tiny amounts of air and steam each time their gauge rocks or jiggles during processing. The sound of the weight rocking or jiggling indicates that the canner is maintaining the recommended pressure and needs no further attention until the load has been processed for the set time. Weighted-gauge canners cannot correct precisely for higher altitudes, and at altitudes above 1,000 feet must be operated at a pressure of 15.

Check dial gauges for accuracy before use each year and replace if they read high by more than 1 pound at 5, 10, or 15 pounds of pressure. Low readings cause overprocessing and may indicate that the accuracy of the gauge is unpredictable. If a gauge is consistently low, you may adjust the processing pressure. For example, if the directions call for 12 pounds of pressure and your dial gauge has tested 1 pound low, you can safely process at 11 pounds of pressure. If the gauge is more than 2 pounds low, it is unpredictable, and it is best to replace it.

Gauges may be checked at most county Extension offices.

Handle gaskets of canner lids carefully and clean them according to the manufacturer's directions. Nicked or dried gaskets will allow steam leaks during pressurization of canners. Gaskets of older canners may need to be lightly coated with vegetable oil once per year, but newer models are pre-lubricated. Check your canner's instructions.

Lid safety fuses are thin metal inserts or rubber plugs designed to relieve excessive pressure from the canner. Do not pick at or scratch fuses while cleaning lids. Use only canners that have Underwriter's Laboratory (UL) approval to ensure their safety.

Replacement gauges and other parts for canners are often available at stores offering canner equipment or from canner manufacturers. Order parts, list canner model number, and describe the parts needed.

Using pressure canners

Follow these steps for successful pressure canning:

1. Put 2 to 3 inches of hot water in the canner. Place filled jars on the rack, using a jar lifter. Fasten canner lid securely.
2. Open petcock or leave weight off vent port. Heat at the highest setting until steam flows from the petcock or vent port.
3. Maintain high heat setting, exhaust steam 10 minutes, and then place weight on vent port or close petcock. The canner will pressurize during the next three to five minutes.
4. Start timing the process when the pressure reading on the dial gauge indicates that the recommended pressure has been reached or when the weighted gauge begins to jiggle or rock.
5. Regulate heat under the canner to maintain a steady pressure at or slightly above the correct gauge pressure. Quick and large pressure varia-

tions during processing may cause unnecessary liquid losses from jars. Weighted gauges on Mirro canners should jiggle about two or three times per minute. On Presto canners, they should rock slowly throughout the process.

6. When processing time is completed, turn off the heat, remove the canner from heat if possible, and let the canner depressurize. *Do not force-cool the canner.* If you cool it with cold running water in a sink or open the vent port before the canner depressurizes by itself, liquid will spurt from jars, causing low liquid levels and jar seal failures. Force-cooling also may warp the canner lid of older model canners, causing steam leaks.

Depressurization of older models should be timed. Standard size heavy-walled canners require about 30 minutes when loaded with pints and 45 minutes with quarts. Newer thin-walled canners cool more rapidly and are equipped with vent locks. These canners are depressurized when their vent lock piston drops to a normal position.

7. After the vent port or petcock has been open for two minutes, unfasten the lid and carefully remove it. Lift the lid away from you so that the steam does not burn your face.
8. Remove jars with a lifter, and place on towel or cooling rack, if desired.

Cooling Jars

Cool the jars at room temperature for 12 to 24 hours. Jars may be cooled on racks or towels to minimize heat damage to counters. The food level and liquid volume of raw-packed jars will be noticeably lower after cooling because air is exhausted during processing and food shrinks. If a jar loses excessive liquid during processing, do not open it to add more liquid. As long as the seal is good, the product is still usable.

Testing Jar Seals

After cooling jars for 12 to 24 hours, remove the screw bands and test seals with one of the following methods:

Method 1: Press the middle of the lid with a finger or thumb. If the lid springs up when you release your finger, the lid is unsealed and reprocessing will be necessary.

Method 2: Tap the lid with the bottom of a teaspoon. If it makes a dull sound, the lid is not sealed. If food is in contact with the underside of the lid, it will also cause a dull sound. If the jar lid is sealed correctly, it will make a ringing, high-pitched sound.

Method 3: Hold the jar at eye level and look across the lid. The lid should be concave (curved down slightly in the center). If center of the lid is either flat or bulging, it may not be sealed.

Reprocessing Unsealed Jars

If a jar fails to seal, remove the lid and check the jar-sealing surface for tiny nicks. If necessary, change the jar, add a new, properly prepared lid, and reprocess within 24 hours using the same processing time.

Another option is to adjust headspace in unsealed jars to 1½ inches and freeze jars and contents instead of reprocessing. However, make sure jars have straight sides. Freezing may crack jars with “shoulders.”

Foods in single unsealed jars could be stored in the refrigerator and consumed within several days.

Storing Canned Foods

If lids are tightly vacuum-sealed on cooled jars, remove screw bands, wash the lid and jar to remove food residue, then rinse and dry jars. Label and date the jars and store them in a clean, cool, dark, dry place. Do not store jars at temperatures above 95°F or near hot pipes, a range, a furnace, in an uninsulated attic, or in direct sunlight. Under these conditions, food will lose quality in a few weeks or months and may spoil. Dampness may corrode metal lids, break seals, and allow recontamination and spoilage.

Accidental freezing of canned foods will not cause spoilage unless jars become unsealed and recontaminated. However, freezing and thawing may soften food. If jars must be stored where they may freeze, wrap them in newspapers, place them in heavy cartons, and cover them with more newspapers and blankets.

Identifying and Handling Spoiled Canned Food

Growth of spoilage bacteria and yeast produces gas which pressurizes the food, swells lids, and breaks jar seals. As each stored jar is selected for use, examine its lid for tightness and vacuum. Lids with concave centers have good seals.

Next, while holding the jar upright at eye level, rotate the jar and examine its outside surface for streaks of dried food originating at the top of the jar. Look at the contents for rising air bubbles and unnatural color.

While opening the jar, smell for unnatural odors and look for spurting

liquid and cottonlike mold growth (white, blue, black, or green) on the top food surface and underside of lid. Do not taste food from a stored jar you discover to have an unsealed lid or that otherwise shows signs of spoilage.

All suspect containers of spoiled low-acid foods should be treated as having produced botulinum toxin and should be handled carefully as follows:

- If the suspect glass jars are unsealed, open, or leaking, they should be detoxified before disposal.
- If the suspect glass jars are sealed, remove lids and detoxify the entire jar, contents, and lids.

Detoxification process

Carefully place the suspect containers and lids on their sides in an 8-quart volume or larger stockpot, pan, or boiling-water canner. Wash your hands thoroughly. Carefully add water to the pot. The water should completely cover the containers with a minimum of a 1 inch of water above the containers. Avoid splashing the water. Place a lid on the pot and heat the water to boiling. Boil 30 minutes to ensure detoxifying the food and all container components. Cool and discard lids and food in the trash or bury in the soil.

Thoroughly clean all counters, containers, and equipment, including can opener, clothing, and hands that may have come in contact with the food or the containers. Discard any sponges or washcloths that were used in the cleanup. Place them in a plastic bag and discard in the trash.

Canned Foods for Special Diets

The cost of commercially canned special diet food often prompts interest in preparing these products at home. Some low-sugar and low-salt foods may be easily and safely canned at home. However, the color, flavor, and texture of these foods may be different than expected and be less acceptable.

Canning without sugar

In canning regular fruits without sugar, it is very important to select fully ripe but firm fruits of the best quality. Prepare these as described for hot packs in FCS3-326, *Selecting, Preparing, and Canning Fruit and Fruit Products*. Juice made from the fruit being canned is best. Blends of unsweetened apple, pineapple, and white grape juice are also good for pouring over solid fruit pieces. Adjust headspaces and lids and use the processing recommendations for regular fruits. Add sugar substitutes, if desired, when serving. Also see FCS3-331, *Preparing and Canning Jams and Jellies*.

Canning without salt (reduced sodium)

To can tomatoes, vegetables, meats, poultry and seafoods, use the procedures given in other publications in the home-canning series, but omit the salt. Consult your county Extension office for these publications: FCS3-327, *Selecting, Preparing, and Canning Tomatoes and Tomato Products*; FCS3-328, *Selecting, Preparing, and Canning Vegetables and Vegetable Products*; and FCS3-329, *Preparing and Canning Poultry, Red Meat, and Fish*. In these products, salt seasons the food but is not necessary to ensure its safety. Add salt substitutes, if desired, when serving.

Glossary

Acid Foods—Foods that contain enough acid to result in a pH of 4.6 or lower. Includes most tomatoes; fermented and pickled vegetables; relishes; jams, jellies, and marmalades; and all fruits except figs. Acid foods may be processed in boiling water.

Ascorbic Acid—The chemical name for vitamin C; commonly used to prevent browning of peeled, light-colored fruits and vegetables.

Blancher—A 6- to 8-quart lidded pot designed with a fitted, perforated basket to hold food in boiling water or with a fitted rack to steam foods. Useful for loosening skins on fruits to be peeled or for heating foods to be hot packed.

Boiling-Water Canner—A large, standard-sized, lidded kettle with jar rack designed for heat-processing 7 quarts or 8 to 9 pints in boiling water.

Botulism—An illness caused by eating toxin produced by growth of *Clostridium botulinum* bacteria in moist, low-acid food containing less than 2 percent oxygen and stored between 40°F and 120°F. Proper heat processing destroys this bacterium in canned food. Freezer temperatures inhibit its growth in frozen food. Low moisture controls its growth in dried food. High oxygen controls its growth in fresh foods.

Canning—A method of preserving food which employs heat processing in airtight, vacuum-sealed containers so that food can be safely stored at normal home temperatures.

Canning Salt—Also called pickling salt. It is regular table salt without the anti-caking or iodine additives.

Citric Acid—A form of acid that can be added to canned foods. It increases the acidity of low-acid foods and may improve their flavor.

Cold Pack—Canning procedure in which jars are filled with raw food. “Raw pack” is the preferred term for describing this practice. “Cold pack” is often used

incorrectly to refer to foods that are open-kettle canned or jars that are heat-processed in boiling water.

Enzymes—Proteins in food which accelerate many flavor, color, texture, and nutritional changes, especially when food is cut, sliced, crushed, bruised, and exposed to air. Proper blanching or hot-packing practices destroy enzymes and improve food quality.

Exhausting—Removal of air from within and around food and from jars and canners. Exhausting or venting of pressure canners is necessary to prevent botulism in low-acid canned foods.

Headspace—The unfilled space above food or liquid in jars that allows for food expansion as jars are heated and for forming vacuums as jars cool.

Heat Processing—Treatment of jars with sufficient heat to enable storing food at normal home temperatures.

Hermetic Seal—An absolutely airtight container seal which prevents reentry of air or microorganisms into packaged foods.

Hot Pack—Heating of raw food in boiling water or steam and filling it hot into jars.

Low-Acid Foods—Foods that contain very little acid and have a pH above 4.6. The acidity in these foods is insufficient to prevent the growth of botulism bacteria. Vegetables, some varieties of tomatoes, figs, all meats, fish, seafoods, and some dairy products are low-acid foods. To control all risks of botulism, jars of these foods must be (1) heat processed in a pressure canner or (2) acidified to a pH of 4.6 or lower before processing in boiling water.

Microorganisms—Independent organisms of microscopic size, including bacteria, yeast, and mold. In a suitable environment, they grow rapidly and may divide or reproduce every 10 to 30 minutes. Therefore, they reach high populations very quickly. Microorganisms are sometimes intentionally added to ferment foods, make antibiotics, and for

other reasons. Undesirable microorganisms cause disease and food spoilage.

Mold—A fungus-type microorganism whose growth on food is usually visible and colorful. Molds may grow on many foods, including acid foods like jams and jellies and canned fruits. Recommended heat processing and sealing practices prevent their growth on these foods.

Mycotoxins—Toxins produced by the growth of some molds on foods.

Open-Kettle Canning—A nonrecommended canning method. Food is heat-processed in a covered kettle, filled while hot into sterile jars, and then sealed. Foods canned this way have low vacuums or too much air, which permits rapid loss of quality in foods. Also, these foods often spoil because they become recontaminated while the jars are being filled.

Pasteurization—Heating food to temperatures high enough to destroy disease-causing microorganisms.

pH—A measure of acidity or alkalinity. Values range from 0 to 14. A food is neutral when its pH is 7.0: lower values are increasingly more acidic; higher values are increasingly more alkaline.

PSIG—Pounds per square inch of pressure as measured by a gauge.

Pressure Canner—A specifically designed metal kettle with a lockable lid used for heat processing low-acid food. These canners have jar racks, one or more safety devices, systems for exhausting air, and a way to measure or control pressure. Canners with 20- to 21-quart capacity are common. The minimum size of canner that should be used has a 16-quart capacity and can hold 7 quart-jars. Use of pressure saucepans with a capacity of less than 16 quarts is not recommended.

Raw Pack—The practice of filling jars with raw, unheated food. Acceptable for canning low-acid foods but allows more rapid quality losses in acid foods that are heat-processed in boiling water. Also called “cold pack.”

Style of Pack—Form of canned food, such as whole, sliced, piece, juice, or sauce. The term may also be used to reveal whether food is filled raw or hot into jars.

Vacuum—A state of negative pressure that reflects how thoroughly air is removed from within a jar of processed food; the higher the vacuum, the less air left in the jar.

Do Your Canned Foods Pass This Quality Test?

OVERALL APPEARANCE

- Good proportion of solid to liquid
- Full pack with proper headspace
- Liquid just covering solid
- Free of air bubbles
- Free of imperfections—stems, cores, seeds
- Good seals
- Practical pack that is done quickly and easily

FRUITS & VEGETABLES

- Pieces uniform in size and shape
- Characteristic, uniform color
- Shape retained—not broken or mushy
- Proper maturity

LIQUID OR SYRUP

- Clear and free from sediment